



CORPORATE PRESENTATION


September 2021

SAFE HARBOR STATEMENT

This presentation contains forward-looking statements. Crinetics cautions you that statements contained in this presentation regarding matters that are not historical facts are forward-looking statements. These statements are based on the company's current beliefs and expectations. Such forward-looking statements include, but are not limited to, statements regarding: the potential benefits of paltusotine for acromegaly patients and patients with carcinoid syndrome; the potential to initiate Phase 3 trials of paltusotine in acromegaly, report data therefrom and the expected timing thereof; the potential for such Phase 3 program to support broad approval of paltusotine for all acromegaly patients who require pharmacotherapy; the potential to initiate a trial of paltusotine in patients with carcinoid syndrome due to NETs and the expected timing thereof; the potential benefits of CRN04894 in patients across multiple indications and the expected timing of the advancement of such program, including the potential to enroll a Phase 1 trial of CRN04894, report data therefrom, and the timing thereof; the potential benefits of CRN04777 in patients with congenital hyperinsulinism and the expected timing of the advancement of such program, including the potential to enroll a Phase 1 trial of CRN04777, report data therefrom, and the timing thereof; the potential for any of our ongoing clinical trials to show safety or efficacy; and our plans to identify and create new drug candidates for additional diseases. In some cases, you can identify forward-looking statements by terms such as "may," "believe," "anticipate," "could," "should," "estimate," "expect," "intend," "plan," "project," "will," "forecast" and similar terms. These statements involve known and unknown risks, uncertainties and other important factors that may cause our actual results, performance or achievements to be materially different from any future results, performance or achievements expressed or implied by the forward-looking statements, including, without limitation: the risk that preliminary results of a clinical trial do not necessarily predict final results and that one or more of the clinical outcomes may materially change as patient enrollment continues, following more comprehensive reviews of the data, and as more patient data become available; potential delays in the commencement, enrollment and completion of clinical trials and the reporting of data therefrom; the FDA or other regulatory agencies may require one or more additional clinical trials of paltusotine or suggest changes to our planned Phase 3 clinical trials prior to and in support of the approval of a New Drug Application or applicable foreign regulatory approval; advancement of paltusotine into a Phase 3 trial in acromegaly or a trial for carcinoid syndrome are dependent on and subject to the receipt of further feedback from the FDA; the COVID-19 pandemic may disrupt our business and that of the third parties on which we depend, including delaying or otherwise disrupting our clinical trials and preclinical studies, manufacturing and supply chain, or impairing employee productivity; our dependence on third parties in connection with product manufacturing, research and preclinical and clinical testing; the success of our clinical trials and nonclinical studies for paltusotine, CRN04894, CRN04777, and our other product candidates; regulatory developments in the United States and foreign countries; unexpected adverse side effects or inadequate efficacy of our product candidates that may limit their development, regulatory approval and/or commercialization; our ability to obtain and maintain intellectual property protection for our product candidates; we may use our capital resources sooner than we expect; and other risks described under the heading "Risk Factors" in documents we file from time to time with the Securities and Exchange Commission. Because forward-looking statements are inherently subject to risks and uncertainties, some of which cannot be predicted or quantified and some of which are beyond our control, you should not rely on these forward-looking statements as predictions of future events. The events and circumstances reflected in our forward-looking statements may not be achieved or occur and actual results could differ materially from those projected in the forward-looking statements. All forward-looking statements are qualified in their entirety by this cautionary statement, which is made under the safe harbor provisions of the Private Securities Litigation Reform Act of 1995 and, except as required by applicable law, we do not plan to publicly update or revise any forward-looking statements contained herein, whether as a result of any new information, future events, changed circumstances or otherwise.


This presentation also contains estimates and other statistical data made by independent parties and by us relating to market size and growth and other data about our industry. This data involves a number of assumptions and limitations, and you are cautioned not to give undue weight to such estimates. In addition, projections, assumptions, and estimates of our future performance and the future performance of the markets in which we operate are necessarily subject to a high degree of uncertainty and risk.

Mission: To build the leading endocrine company that consistently pioneers new therapeutics to help patients better control their disease and improve their daily lives




Strategy: Drugs Built from Scratch for Purpose

We aim to discover, develop and commercialize drugs for endocrine indications with:



Endocrinology Development Strategy: Focus on Hormone Levels from Preclinical to Approval



Pipeline Targets Indications with Unmet Need



Well established biology has the potential to allow for the use of biomarkers as registration endpoints

PROGRAM	Development Stage				Registrational Endpoint	Prevalence	
	Preclin	Phase 1	Phase 2	Phase 3		US Total	Global Range per 100,000
Paltusotine (SST2 agonist)		Pharmacologic POC					
Acromegaly					IGF-1 Levels	26K	2.8 – 13.0
Carcinoid Syndrome					Diarrhea & Flushing	33K	3.7 – 9.7
Nonfunctional NETs					Anti-tumor activity	138K	17 – 46
CRN04894 (ACTH antagonist)							
Cushing's Disease					Cortisol Levels	10K	2.5 – 3.8
Congenital Adrenal Hyperplasia (CAH)					Adrenal Androgens/ Glucocorticoid use	27K	6.7 – 10.0
CRN04777 (SST5 agonist)							
Congenital Hyperinsulinism					Glucose Levels/ Hypoglycemic Events	1.5-2K	0.64 – 1.3
Syndromic Hyperinsulinism					Glucose Levels/ Hypoglycemic Events	2K	Variable


We continue to invest in ongoing discovery of new molecules targeting new indications with our in-house discovery team who are actively pursuing multiple programs in parallel. Indications include hyperparathyroidism, nonfunctional pituitary adenomas, polycystic kidney disease and more.

Endocrinology Invented Biomarkers!



Pharmacologic POC in Phase 1 is Our Goal



Phase 1 SAD Data:
Suppression of GH



Phase 1 MAD Data:
Suppression of IGF-1



PALTUSOTINE: AN INVESTIGATIONAL, POTENTIAL FIRST-IN-CLASS, ORAL NONPEPTIDE SST2 AGONIST

Acromegaly

Carcinoid syndrome

Nonfunctional neuroendocrine tumors




Acromegaly and NETS are Currently Treated with Injected SST2 Peptide Agonists

Acromegaly

- Caused by benign pituitary tumor that secretes excess growth hormone (GH)
- Excess GH causes excess secretion of insulin-like growth factor-1 (IGF-1)

Results in:

1. Bone and cartilage overgrowth
2. Organ enlargement
3. Changes in glucose and lipid metabolism
4. Abnormal growth of hands and feet
5. Alteration of facial features




US Prevalence: **26,000**

SST2 agonist

Neuroendocrine Tumors (NETs)

- NETs arise from aberrant enteroendocrine cells in GI, pancreas or lungs
- In ~19% of cases, tumors are associated with excess secretion of serotonin and other hormones resulting in carcinoid syndrome
- Majority of tumors are “nonfunctional” and not associated with secretory syndrome
- Patients with grade 1 and 2 NETs and distant metastases have a 5-year survival ranging from 30-70%



US Prevalence	
Carcinoid Syndrome:	33,000
Nonfunctional tumors:	138,000
Total	171,000





~\$3B Market Despite Limitations of Current Therapies




Sandostatin (octreotide)	Somatuline (lanreotide)	Somavert (pegvisomant)	MYCAPSSA (oral octreotide)
 NOVARTIS	 IPSEN Innovation for patient care	 Pfizer	 CHIASMA
\$1.4B	\$1.4B	\$277M	\$1M
Monthly intramuscular 5-mL vial; 1½" 19-gauge needle	Monthly deep subcutaneous .2-5ml; 18-gauge needle	Daily injections 1 ml; 28 – 31- gauge needle not supplied	<u>Twice</u> daily oral capsule
Painful injections. Injection site reactions Inconvenient monthly visits to physician's office interrupts normal life Limited efficacy , as many patients experience return of symptoms near end of month	Painful injections. Injection site reactions Inconvenient monthly visits to physician's office interrupts normal life Limited efficacy , as many patients experience return of symptoms near end of month	Inconvenient. Daily dose kits require refrigeration. Patients often must buy a second refrigerator for storage, making travel difficult.	Food effect. Cannot be taken <1 hour before eating or <2 hours after eating ⁽¹⁾ Limited efficacy , as 42% of pivotal study patients did not maintain IGF-1 biochemical response after switching to MYCAPSSA from injectables ⁽¹⁾ Multiple drug-drug interactions ⁽¹⁾ Cold chain distribution ⁽¹⁾
Approval date: 1988, 1998(LAR)	Approval date: 2007	Approval date: 2003	Approval date: 2020

⁽¹⁾MYCAPSSA Label



Paltusotine (CRN00808)	
Oral Bioavailability	70%
Observed Half Life	42-50 hr.

Paltusotine was Designed to Avoid Drug-Drug Interactions and Be Easy to Prescribe



Unchanged paltusotine was the primary circulating species (>75%)

Excretion Summary	Fraction in Urine (%)	Fraction in Feces (%)	Total Recovery (%)
Mean (%CV)	3.9 (29.9)	90 (2.5)	94 (1.1)

Paltusotine has not shown clinically meaningful inhibition or induction of drug metabolizing enzymes

- Dose adjustment may not be required for co-administered drugs



Low risk for drug interactions from inhibition of drug-metabolizing enzymes by co-administered drugs

- Paltusotine is eliminated via the liver
- Multiple potential metabolic pathways
 - Glucuronidation and CYP-dependent metabolism
 - No single pathway is dominant

No excipients that may influence permeability of co-administered drugs

Phase 2 ACROBAT Edge Study Met Primary Endpoint


Hormone suppression maintained after switching from injected SOC to oral paltusotine




Data presented are median (Interquartile Range [IQR]: 25th percentile, 75th percentile) from the EDGE study's primary analysis population. EoT = End of Treatment defined as Week 13 (Visit 14) or last on treatment value carried forward (LOCF). Wks after WD is defined as Week 17 or result at least 22 days after last dose. Note: p-values are based on non-parametric Wilcoxon Sign Rank test of whether the median change is different from zero.

Evidence of Dose Response from Phase 2 Program

Results from Switching to Paltusotine: Change in IGF-1 from Baseline to Steady State at Indicated Dose



Magnitude of Paltusotine Activity: Change in IGF-1 from Steady State to 4 Weeks After Withdrawal



Post-hoc analysis of data from EDGE primary analysis population and EVOLVE patients

Steady state IGF-1 at the indicated dose: Patients were on the indicated dose for at least 12 days. Data prior to Week 7 were excluded because of insufficient washout of depot injection during this window. Data are shown from Week 7, Week 10, and Week 13.

Data presented are median $\pm$ IQR. EoT = End of Treatment defined as Week 13 (Visit 14) or last on treatment value carried forward (LOCF).

WD = Withdrawal, where 2 Wks after WD is defined as Week 15 or result within 7-21 days after last dose and 4 Wks after WD is defined as Week 17 or result at least 22 days after last dose. One subject is missing 4 Weeks after withdrawal observation.

Octreotide and lanreotide concentrations were measured 17 weeks after depot dose (W13 of the treatment period). Octreotide was completely washed out. Lanreotide concentrations were $>75\%$ reduced from baseline.



Phase 3 Program: Designed to Support Potential for Broad First-Line Medical Therapy

Two double-blind, placebo-controlled studies planned to support broad labeling in the U.S. and Europe for use in all acromegaly patients who require pharmacotherapy

PATHFINDER-1: Switching from SOC

Evaluate safety and efficacy of paltusotine in acromegaly patients switching from injectable octreotide or lanreotide depots, who are currently *biochemically controlled*

PATHFINDER-2: Untreated Patients

Evaluate safety and efficacy of paltusotine in untreated acromegaly patients who are *biochemically uncontrolled*

Anticipated Paltusotine Milestones

- 1 Initiate PATHFNDR-1: switching from SOC ✓
- 2 Initiate PATHFNDR-2: use in untreated patients (anticipated in 2H 2021)
- 3 Initiation of Phase 2 NETs trial in carcinoid syndrome (end of 2021)
- 4 Report topline data from PATHFNDR-1 & 2 trials (expected in 2023)


CRN04894: AN INVESTIGATIONAL, POTENTIAL FIRST-IN-CLASS, ORAL NONPEPTIDE ACTH ANTAGONIST

Congenital adrenal hyperplasia (CAH)

Cushing's disease (CD)


Other conditions of ACTH excess

The Hypothalamic-Pituitary-Adrenal (HPA) Axis is the Body's Emergency Response System for Stress




Hypothesis: An Oral, Selective ACTH Antagonist Will Have Utility in Treating Diseases of ACTH

Normal




ACTH Range = 5 – 60 pg/mL ^{1,4,5}

Cushing's Disease (CD)



ACTH Range = 20-200 pg/mL ¹

Congenital Adrenal Hyperplasia (CAH)




ACTH Range = 150-500 pg/mL ^{2,3}

Cause	ACTH-secreting pituitary tumor	Inability to produce cortisol leads to loss of negative feedback & excess ACTH
US Prevalence (global incidence per 100,000)	10k (2.5-3.8)	27k (6.7-10.0)
Symptoms	Central obesity and round face; Dorsal and supraclavicular fat pads; Hypertension; Stretch marks; Bone loss; Hyperglycemia; Psychiatric disturbances	Adrenal insufficiency; Infertility; Hirsutism; Short stature; Precocious puberty; Adrenal rest tumors


¹ Raff et al. Compr Physiol 2015, ² Petersen Acta Paediatr Scand 1981, ³ NBIX ENDO Online 2020 presentation, ⁴ Oster et al., Endocrine Reviews 2017, ⁵ UpToDate Reference, ⁶ Oelkers et al, JCEM 1988, ⁶ Alia et. al Clinical Endocrinology 2006

CRN04894: First-in-Class ACTH Antagonist for ACTH Driven Diseases



There Are No ACTH Receptor Blocking Agents Available to Treat ACTH Driven Diseases

All currently approved agents and agents in development act upstream or downstream of ACTH




References: Felders et al. Lancet Diab Endo 7:300-12, 2019. Castinetti JCEM 99: 1623-1639, 2014. Castinetti JCEM 106: 2114-2123, 2021.

CRN04894 is the Only ACTH Antagonist in Clinical Development

CRN04894 was carefully crafted by Crinetics in-house discovery team


CRN04894 is a potent ($K_b = 0.4$ nM) competitive antagonist of ACTH signaling



Mechanism of action

- Designed to compete with ACTH for a common binding site in order to block the ACTH-induced signaling.
- Relative affinity and concentration of CRN04894 and ACTH potentially determine balance of occupancy (competitive antagonism).

Acute suppression of ACTH-induced corticosterone observed in rats



Experiment designed to mimic disease:

- CRN04894 orally administered
- Administer IV bolus of ACTH after 60 minutes
- Marked suppression of ACTH with increasing doses of CRN04894
- Analogous ACTH challenge in Phase 1 POC


CRN04894 SAD Study Design to Establish Pharmacologic Proof-of-Concept

Follows Crinetics' core endocrine strategy of using hormonal biomarkers to drive development

Study Goals

- Evaluate safety [10-80 mg]
- Evaluate pharmacokinetics: oral absorption, dose-proportional exposure, half-life [10-80 mg]
- Evaluate dose-response & PK/PD on basal cortisol [10-80 mg]
- Evaluate dose-response & PK/PD using supra-pathophysiologic ACTH challenge (250 mcg) [10-80 mg]
- Evaluate cortisol suppression with selected dose in response to disease-relevant ACTH challenge (1 mcg) [80 mg only]


ACTH Challenge Test



Proof of concept: dose dependent suppression of basal cortisol and ACTH-stimulated cortisol with CRN04894

PK Results: CRN04894 Showed Oral Bioavailability With Dose-Proportional Exposure


Half-life ~24 hour and t_{max} ~1 hour



Data shown are mean±SEM

CRN04894 Rapidly Reduced Basal Cortisol Output from Adrenal Glands



Acute reduction of basal cortisol (56% @ 80 mg) 2 hours after administration of CRN04894



Data shown are mean±SEM
^a Full suppression of cortisol production assumes no more cortisol is produced at time of CRN04894 dose and cortisol half-life is 66 ±18 min from McKay LI, Cidlowski JA. Pharmacokinetics of Corticosteroids. In: Kufe DW, Pollock RE, Weichselbaum RR, et al., editors. Holland-Frei Cancer Medicine. 6th edition. Hamilton (ON); 2003

Dose-Dependent Suppression of Cortisol Observed Following Supra-Pathophysiologic ACTH Challenge (250 mcg)


CRN04894 resulted in strong cortisol suppression (41% @ 80 mg) despite anticipated ACTH exposure orders of magnitude higher than disease states



Data shown are mean $\pm$ SEM

Clinically Meaningful Cortisol Suppression Observed in Response to Disease-relevant ACTH Challenge (1 mcg)

CRN04894 maintains normal cortisol levels for these subjects in face of disease-relevant ACTH (1 mcg) challenge



Data shown are mean±SEM

Conclusions from CRN04894 SAD Results

Objectives

- Safety and tolerability
- Drug-like Pharmacokinetics
- PK/PD for suppression of ACTH-induced adrenal activity

Generally safe and well tolerated at single doses from 10 to 80 mg



Achieved targeted pharmacokinetic profile

- Rapidly absorbed after oral administration (t_{max} ~1 hr)
- Dose proportional exposure from 10 to 80 mg
- Favorable half-life of ~24 hours



Demonstrated pharmacologic proof-of-concept for ACTH antagonism

- Strong suppression of basal cortisol (56%)
- Dose-dependent, strong cortisol suppression (41%) following supra-pathophysiologic ACTH (250 mcg) challenge
- Maintains normal cortisol levels for the Phase 1 participants in face of disease-relevant ACTH (1 mcg) challenge



Recent and Anticipated CRN04894 Milestones

- 1 Open US IND (complete) ✓
- 2 Initiate Phase 1 FIH healthy volunteer POC study ✓
- 3 Report Phase 1 SAD data (reported August 2021) ✓
- 4 Report Phase 1 MAD data (expected 1Q 2022)

CRN04777: AN INVESTIGATIONAL, POTENTIAL FIRST-IN-CLASS, ORAL NONPEPTIDE SST5 AGONIST


Congenital hyperinsulinism (HI)



Congenital Hyperinsulinism Results in Life Threatening Recurrent Hypoglycemia

Congenital HI patients secrete insulin even when blood sugar is low

Congenital HI is a devastating rare disease




- Untreated hypoglycemia can result in neurodevelopment disorders and death
- Early identification and continuous intensive glucose management are critical
- Current treatment paradigms place high burden of care on families with all too frequent suboptimal outcomes
- Six Global Centers of Excellence named for treatment of patients with HI
- Robust global patient advocacy such as Congenital Hyperinsulinism International (www.congenitalhi.org)



Serious Unmet Medical Needs in Congenital HI

Intensive 24h-glucose management (monitoring, feeding, glucose tube)

Current Standard of Care for Congenital HI



Patient & Parent Goals


- ✓ Avoid hypoglycemia and its consequences including neurological damage
- ✓ Safely sleep through the night
- ✓ Avoid pancreatectomy
- ✓ Eliminate feeding tubes
- ✓ Reduce injections and glucose sticks
- ✓ Medical management until HI resolves
- ✓ Be a kid not a patient





Congenital HI Patient Care is a High Burden on Healthcare Systems

Healthcare utilization by a baby girl with Congenital HI




Current Challenges

- Variable time to diagnosis
- Constant dextrose infusion to maintain normal blood sugar levels
- Surgical removal of all or part of the pancreas – Or
- No surgical options
- Ineffective diazoxide treatment with multiple untoward effects

As a result:

- Hypoglycemic crises warranting repeat need for emergency services (can include seizure, loss of consciousness and death)
- Frequent and multi-day inpatient hospital stays
- Long-term consequences including neurodevelopmental impairment


Cost of Illness Estimate from the UK², £ 3,408,398 (\$4,630,939): first 11 years of life




Patients unresponsive to first-line drug therapy (diazoxide) represented the greatest driver of costs

1. Claims data on file, 2013-2018 2. Eljamel, S et al The burden of congenital hyperinsulinism in the United Kingdom: a cost of illness study 2018

CRN04777: First-in-class Oral SST5 Agonist with Potential to be Broadly Effective in HI




SST5 Inhibits Insulin Secretion Downstream of all Known HI Causing Mutations




Syndromic hyperinsulinisms (e.g. those associated with Beckwith-Wiedemann syndrome, Sotos syndrome, Kabuki syndrome, and Turner syndrome) may also respond to SST5 agonism


Proof of Mechanism Achieved in Animal Models and Patient Islets



CRN04777 suppressed sulfonylurea (SU)-induced insulin secretion and reversed hypoglycemia in rats



CRNX SST5 agonist suppressed insulin from islets isolated from patient with Beckwith-Wiedemann Syndrome



Islet data was obtained using another Crinetics SST5 agonist candidate before CRN04777 had been selected for development

CRN04777 SAD Study Design to Evaluate Pharmacologic Proof-of-Concept

Follows Crinetics' core endocrine strategy of using hormonal biomarkers to drive development

Study Goals

- Evaluate safety [0.5-120 mg]
- Evaluate pharmacokinetics: oral absorption, dose-proportional exposure, half-life [0.5-120 mg]
- Evaluate dose response and PK/PD on pre- and post-stimulated glucose and insulin in an IVGTT [0.5-120 mg]
- Evaluate dose response and PK/PD on reduction/reversal of sulfonylurea-induced insulin secretion (pharmacologic model of disease) [30-60 mg]

Pharmacodynamic Assessments


1. Intravenous Glucose Tolerance Test (IVGTT)
2. Sulfonylurea (SU) Challenge

Proof-of-Concept

- Dose dependent suppression of glucose- or sulfonylurea-induced insulin secretion with CRN04777

PK Results: CRN04777 Showed Oral Bioavailability with Dose-Proportional Exposure

Half-life ~40 hours and t_{max} ~1-2 hours at efficacious doses



Data shown are mean $\pm$ SEM; LLOQ = lower limit of quantitation

All doses n=6; except n=12 for 60 mg which was evaluated in both IVGTT and sulfonylurea challenge

*When 120 mg was administered within 30 minutes of a standard adult high fat breakfast, a significant reduction in exposure was observed. Evaluation of pediatric relevant meals pending.


CRN04777 SAD Study Designed to Evaluate Pharmacologic Proof-of-Concept

Follows Crinetics' core endocrine strategy of using hormonal biomarkers to drive development

Study Goals

- Evaluate safety [0.5-120 mg]
- Evaluate pharmacokinetics: oral absorption, dose-proportional exposure, half-life [0.5-120 mg]
- **Evaluate dose response and PK/PD on pre- and post-stimulated glucose and insulin in an IVGTT [0.5-120 mg]**
- Evaluate dose response and PK/PD on reduction/reversal of sulfonylurea-induced insulin secretion (pharmacologic model of disease) [30-60 mg]

1. Intravenous Glucose Tolerance Test (IVGTT)





Healthy volunteers received single oral dose of CRN04777
(n = 8, 6 active/ 2 placebo in each cohort)

CRN04777 Dose-Dependently Suppressed Glucose Stimulated Insulin Secretion

CRN04777 dose-dependently reduced insulin secretion stimulated by bolus IV glucose (IVGTT)...

...and reduced insulin secretion reduces glucose uptake by tissues resulting in prolonged elevation of plasma glucose



Data shown are mean±SEM
N=6 CRN04777 treated per dose; N=14 placebo
IVGTT=intravenous glucose tolerance test; PBO=placebo


CRN04777 SAD Study Designed to Evaluate Pharmacologic Proof-of-Concept

Follows Crinetics' core endocrine strategy of using hormonal biomarkers to drive development

Study Goals

- Evaluate safety [0.5-120 mg]
- Evaluate pharmacokinetics: oral absorption, dose-proportional exposure, half-life [0.5-120 mg]
- Evaluate dose response and PK/PD on pre- and post-stimulated glucose and insulin in an IVGTT [0.5-120 mg]
- **Evaluate dose response and PK/PD on reduction/reversal of sulfonylurea-induced insulin secretion (pharmacologic model of disease) [30-60 mg]**

2. Sulfonylurea (SU) Challenge









Healthy volunteers received single oral dose of CRN04777
(n = 9, 6 active / 3 placebo in each cohort)

CRN04777 Reversed Sulfonylurea-Induced Hyperinsulinism in a Pharmacologic Model of Congenital HI

CRN04777 eliminated the need for IV glucose support by inhibiting insulin secretion

Glucose Infusion Rate (GIR) – Increases in Proportion to Insulin Secretion



More detailed data (insulin, C-peptide) to be presented at future medical conferences

Solid line: mean value; shaded area: SEM

Conclusions from CRN04777 SAD Results

Objectives

- Safety and tolerability
- Drug-like pharmacokinetics
- PK/PD for suppression of insulin secretion

Safe and well tolerated at single doses from 0.5-120 mg



Achieved targeted pharmacokinetic profile

- Rapidly absorbed after oral administration ($t_{\max}$ ~1-2 hrs)
- Favorable half-life of ~40 hours observed
- Dose-proportional exposure from 0.5-120 mg



Demonstrated pharmacologic proof-of-concept for SST5 agonism

- Dose-dependent reduction in glucose-induced insulin secretion achieved in an intravenous glucose tolerance test
- Dose-dependent reversal of sulfonylurea-induced insulin secretion achieved in a pharmacologic model of hyperinsulinism



Recent and Anticipated CRN04777 Milestones

- 1 US Rare Pediatric Disease and EU Orphan Drug Designations (received; CRNX may be eligible for priority review voucher in the US) ✓
- 2 Initiate Phase 1 FIH healthy volunteer POC study ✓
- 3 Report Phase 1 SAD data (reported September 2021) ✓
- 4 Report Phase 1 MAD data (expected 1Q 2022)

Achieved 2021 Goal of Three Programs with Clinical Proof-of-Concept

	1H21	2H21	1H22	2H22
Paltusotine SST2 Agonist for Acromegaly & NETs POC Achieved	Initiate PATHFND-1 	Initiate PATHFND-2	PATHFND-1 and PATHFND-2 Ongoing	
		Initiate P2 NETs Trial in Carcinoid Syndrome	Carcinoid Syndrome Phase 2 Ongoing	
CRN04894 ACTH Antagonist for Cushing's Disease & CAH POC Achieved	Initiate Phase 1 	Phase 1 POC SAD Data 	Phase 1 MAD Data(Q1)	Initiate Phase 2 in Patients
CRN04777 SST5 Agonist for Congenital HI POC Achieved	Initiate Phase 1 	Phase 1 POC SAD Data 	Phase 1 MAD Data(Q1)	Initiate Phase 2 in Congenital HI Patients

Executed development strategy with three NCEs: paltusotine, '4894, and '4777

APPENDICES

Key Patent Families Anchor a Robust IP Portfolio

Patent Family Subject Matter	Patent Status	Priority Date	Estimated Expiration
Paltusotine Portfolio			
Composition of Matter	Granted in: US, AU, IN Pending in: foreign jurisdictions representing >96% of pharmaceutical markets	July 2016	July 2037
HCl Salt and its Polymorph Form	Granted in: U.S. Pending in: foreign jurisdictions representing >96% of pharmaceutical markets	January 2018	January 2039
New Formulation	Pending in: PCT, U.S., TW, AR, VE	September 2020	September 2041
Acromegaly Treatment Methods	Pending in: U.S	May 2021	May 2042
CRN04894 Portfolio			
Composition of Matter	Granted in: U.S. Pending in: foreign jurisdictions representing >96% of pharmaceutical markets	June 2018	June 2039
Treatment Methods	Pending in: U.S	March 2021	March 2042
CRN04777 Portfolio			
Composition of Matter	Pending in: PCT, U.S., TW, AR, VE	Aug 2019	Aug 2040
Polymorph Form	Pending in: U.S	February 2021	February 2042
Treatment Methods	Pending in: U.S	February 2021	February 2042

Board of Directors

Wendell Wierenga, PhD Chairman (Former EVP R&D, Santarus)



Scott Struthers, PhD Founder & CEO



Matt Fust, MBA Former CFO, Onyx



Weston Nichols, PhD Founder & Managing Partner, Lynx1 Capital



Stephanie Okey, MS Former SVP, Genzyme Corporation



Camille Bedrosian, MD Chief Medical Officer, Ultragenyx Pharmaceutical



Scientific Advisory Board

David Clemmons, MD

Professor of Medicine at UNC, Chapel Hill



Anne Klibanski, MD

President & CEO, Partners Healthcare
Former Chief of Neuroendocrine Unit at
MGH & Professor of Medicine at Harvard



MASSACHUSETTS
GENERAL HOSPITAL



HARVARD
MEDICAL SCHOOL



Philip Harris, FRCP, PhD

Chief Medical Officer,
Isotopen Technologien München

